

# The Medical Letter<sup>®</sup>

on Drugs and Therapeutics

## Comparison Chart: **ANTIVIRAL DRUGS FOR INFLUENZA**

Updated November 13, 2022

**The Medical Letter<sup>®</sup>**  
**Because the source matters.**

**To support more content like this,  
please consider making a donation\* or becoming a subscriber.**

The Medical Letter is a nonprofit organization<sup>†</sup> that relies solely on subscription fees and donations to support our mission of providing objective, practical, and timely information on drugs and therapeutics.

Our work relies on support from people like you who value credible, unbiased drug information that is free of any commercial interest.

**For more information click: [Subscriptions](#), [Site Licenses](#), [Reprints](#) or call Customer Service at: 800-211-2769**

<sup>†</sup>The Medical Letter is a nonprofit, tax-exempt organization under Section 501(C)(3) of the Internal Revenue Code. Donations are tax-deductible as allowed by law.

## ANTIVIRAL DRUGS FOR INFLUENZA FOR 2022-2023

DRUG	FORMULATIONS	ROUTE	TREATMENT	CHEMOPROPHYLAXIS	COST <sup>1</sup>
<b>NEURAMINIDASE INHIBITORS</b>					
<b>Oseltamivir</b> – generic <i>Tamiflu</i> (Genentech)	30, 45, 75 mg caps; 6 mg/mL oral susp	Oral NG tube	<ul style="list-style-type: none"> <li>FDA-approved for treatment of acute uncomplicated influenza in patients ≥2 weeks old</li> <li>Preferred for treatment of influenza in pregnant women, hospitalized patients, and outpatients with severe, complicated, or progressive illness</li> </ul>	<ul style="list-style-type: none"> <li>FDA-approved for chemoprophylaxis of influenza in patients ≥1 year old</li> </ul>	\$32.00 151.90
<b>Peramivir</b> – <i>Rapivab</i> (BioCryst)	200 mg/20 mL single-use vials	IV	<ul style="list-style-type: none"> <li>FDA-approved for treatment of acute uncomplicated influenza in otherwise healthy patients ≥6 months old</li> <li>Not recommended for treatment of severe influenza (can be considered for patients who cannot tolerate oseltamivir)</li> </ul>	<ul style="list-style-type: none"> <li>Not FDA-approved for chemoprophylaxis</li> </ul>	950.00
<b>Zanamivir</b> – <i>Relenza</i> (GSK)	5 mg blisters of powder for inhalation	Inhalation	<ul style="list-style-type: none"> <li>FDA-approved for treatment of acute uncomplicated influenza in patients ≥7 years old</li> <li>Not recommended for patients with underlying airway disease</li> <li>Not recommended for treatment of severe influenza</li> </ul>	<ul style="list-style-type: none"> <li>FDA-approved for chemoprophylaxis of influenza in patients ≥5 years old</li> </ul>	59.00
<b>POLYMERASE ACIDIC (PA) ENDONUCLEASE INHIBITOR</b>					
<b>Baloxavir marboxil</b> – <i>Xofluza</i> (Genentech)	40, 80 mg tabs; 40 mg/20 mL oral susp	Oral	<ul style="list-style-type: none"> <li>FDA-approved for treatment of acute uncomplicated influenza in otherwise healthy patients ≥5 years old or in patients ≥12 years old who are at high risk of influenza-related complications</li> <li>No data in patients with severe influenza</li> <li>Not recommended for use in severely immunosuppressed persons or pregnant women</li> </ul>	<ul style="list-style-type: none"> <li>FDA-approved for post-exposure chemoprophylaxis in patients ≥5 years old</li> </ul>	154.50

### RECOMMENDATIONS FOR ANTIVIRAL TREATMENT<sup>2,3</sup>

- Antiviral treatment should be started as soon as possible; it is most effective when started within 48 hours after illness onset
- Antiviral treatment is recommended for patients with suspected or confirmed influenza who are hospitalized, have severe, complicated, or progressive illness, or are at increased risk for complications, even if it is started >48 hours after illness onset
- Antiviral treatment can be considered for otherwise healthy symptomatic outpatients with suspected or confirmed influenza who are not at increased risk for influenza complications if it can be started within 48 hours after illness onset

### PREGNANCY AND LACTATION

- Oseltamivir and zanamivir appear to be safe for use during pregnancy
- Oseltamivir is preferred for treatment of women who are pregnant, ≤2 weeks postpartum, or breastfeeding
- No data are available on use of baloxavir in pregnant or breastfeeding women
- Zanamivir may be preferred for chemoprophylaxis in pregnant women because of its limited systemic absorption, but oseltamivir is a reasonable alternative, especially in women who are at increased risk for respiratory problems

## ANTIVIRAL DRUGS FOR INFLUENZA FOR 2022-2023 (continued)

DRUG	USUAL TREATMENT DOSAGE	USUAL CHEMOPROPHYLAXIS DOSAGE
<b>OSELTAMIVIR (TAMIFLU)</b>		
<b>ADULT</b>	75 mg PO bid x 5 days	75 mg PO once/day x 7 days
<b>PEDIATRIC</b>	<b>&lt;2 wks old:</b> 3 mg/kg PO bid x 5 days (CDC) <sup>4</sup> <b>≥2 wks-&lt;1 yr:</b> 3 mg/kg PO bid x 5 days (9-11 months: AAP recommends 3.5 mg/kg) <sup>5</sup> <b>1-12 yrs:</b> 30-75 mg PO bid x 5 days (≤15 kg: 30 mg; >15-23 kg: 45 mg; >23-40 kg: 60 mg; >40 kg: 75 mg) <b>≥13 yrs:</b> 75 mg PO bid x 5 days	<b>&lt;3 mos:</b> not recommended <b>3 mos-1 yr:</b> 3 mg/kg PO once/day x 7 days (CDC & AAP) <sup>6</sup> <b>1-12 yrs:</b> 30-75 mg PO once/day x 7 days (≤15 kg: 30 mg; >15-23 kg: 45 mg; >23-40 kg: 60 mg; >40 kg: 75 mg) <b>≥13 yrs:</b> 75 mg PO once/day x 7 days
<b>RENAL</b>	<b>Adults and children &gt;40 kg (CDC):</b> <ul style="list-style-type: none"> <li>▪ CrCl 31-60 mL/min: 30 mg bid</li> <li>▪ CrCl 11-30 mL/min: 30 mg once/day</li> <li>▪ HD: 30 mg after every HD (may be started immediately if influenza symptoms develop between HD sessions)</li> <li>▪ CAPD: 30 mg once immediately</li> <li>▪ ESRD (not on HD): not recommended</li> </ul>	<b>Adults and children &gt;40 kg (CDC):</b> <ul style="list-style-type: none"> <li>▪ CrCl 31-60 mL/min: 30 mg once/day</li> <li>▪ CrCl 11-30 mL/min: 30 mg every other day</li> <li>▪ HD: 30 mg after every other HD (initial dose can be given before start of HD)</li> <li>▪ CAPD: 30 mg immediately and then once/week after exchange</li> <li>▪ ESRD (not on HD): not recommended</li> </ul>
<b>PERAMIVIR (RAPIVAB)</b>		
<b>ADULT</b>	600 mg IV over 15-30 minutes once	Not FDA-approved for chemoprophylaxis
<b>PEDIATRIC</b>	<b>6 months-12 yrs:</b> 12 mg/kg (max 600 mg) IV over 15-30 mins once <b>≥13 yrs:</b> 600 mg IV over 15-30 mins once	Not FDA-approved for chemoprophylaxis
<b>RENAL</b>	<b>2-12 yrs:</b> <ul style="list-style-type: none"> <li>▪ CrCl 30-49 mL/min: 4 mg/kg IV once</li> <li>▪ CrCl 10-29 mL/min: 2 mg/kg IV once</li> </ul> <b>≥13 yrs:</b> <ul style="list-style-type: none"> <li>▪ CrCl 30-49 mL/min: 200 mg IV once</li> <li>▪ CrCl 10-29 mL/min: 100 mg IV once</li> <li>▪ HD: administer dose after HD (based on CrCl)</li> </ul>	
<b>ZANAMIVIR (RELENZA)</b>		
<b>ADULT</b>	2 inhalations bid x 5 days	2 inhalations once/day x 7 days
<b>PEDIATRIC</b>	≥7 yrs: 2 inhalations bid x 5 days	≥5 yrs: 2 inhalations once/day x 7 days
<b>BALOXAVIR (XOFLUZA)</b>		
<b>ADULT</b>	<b>&lt;80 kg:</b> 40 mg PO once <b>≥80 kg:</b> 80 mg PO once	<b>&lt;80 kg:</b> 40 mg PO once <b>≥80 kg:</b> 80 mg PO once
<b>PEDIATRIC</b>	<b>≥5 yrs and &lt;80 kg:</b> 40 mg PO once <b>≥5 yrs and ≥80 kg:</b> 80 mg PO once	<b>≥5 yrs and &lt;80 kg:</b> 40 mg PO once <b>≥5 yrs and ≥80 kg:</b> 80 mg PO once

### DURATION OF THERAPY

#### **TREATMENT:**

- Oseltamivir or zanamivir should be given for 5 days; peramivir and baloxavir are given as single doses
- In hospitalized, critically ill, or immunocompromised patients, a longer treatment course of oseltamivir (e.g., 10 days) is often used. IV peramivir for at least 5 days may be considered for hospitalized, critically ill patients, or patients who cannot tolerate or absorb oral or enterically administered oseltamivir because of gastric stasis, malabsorption, or GI bleeding

#### **CHEMOPROPHYLAXIS:**

- Oseltamivir or zanamivir should be continued for 7 days after the last known exposure
- For institutional outbreaks, the CDC recommends that chemoprophylaxis be given for at least 2 weeks and continued for up to 1 week after the end of the outbreak

### ADMINISTRATION

- Taking oseltamivir with food may minimize GI adverse effects
- Oseltamivir capsules can be opened and the contents mixed in a thick sweetened liquid to mask the bitter taste and consumed immediately
- Oseltamivir can be given by oro/nasogastric tube to patients who are unable to swallow
- Baloxavir suspension must be used within 10 hours after reconstitution

AAP = American Academy of Pediatrics; CAPD = continuous ambulatory peritoneal dialysis; CDC = Centers for Disease Control; ESRD = end-stage renal disease; HD = hemodialysis

## ANTIVIRAL DRUGS FOR INFLUENZA FOR 2022-2023 (continued)

### SOME ADVERSE EFFECTS

#### NEURAMINIDASE INHIBITORS AND BALOXAVIR:

- Neuropsychiatric events have been reported, but a causal relationship has not been established
- Neuropsychiatric dysfunction can be a complication of influenza itself

#### OSELTAMIVIR:

- Nausea, vomiting, headache

#### PERAMIVIR:

- Diarrhea, neutropenia

#### ZANAMIVIR:

- Diarrhea, nausea, sinusitis, fever, arthralgia
- Bronchospasm; not recommended in patients with underlying airway disease
- Contains lactose; contraindicated in patients with a history of milk protein allergy

#### BALOXAVIR:

- Nausea and vomiting; incidence appears to be lower than with oseltamivir

### DRUG INTERACTIONS

#### With Intranasal Live-Attenuated Influenza Vaccine (*Flumist Quadrivalent*):

- Antivirals could inhibit replication of vaccine virus and reduce vaccine efficacy
- Avoid **oseltamivir** or **zanamivir** within 48 hours before, **peramivir** within 5 days before, or **baloxavir** within 17 days before vaccine administration
- Revaccination with an inactivated or a recombinant influenza vaccine is recommended in persons who receive any one of these antiviral drugs during these specified times and through 2 weeks after receiving the intranasal influenza vaccine

#### With Polyvalent Cations:

- Coadministration of dairy products, beverages, antacids, laxatives, multivitamins, or other products containing polyvalent cations (calcium, aluminum, iron, magnesium, selenium, or zinc) can reduce serum concentrations of **baloxavir** and should be avoided.

### ACTIVITY/RESISTANCE

- ▶ Neuraminidase inhibitors and baloxavir have activity against influenza A and B viruses
- ▶ Over 99% of recently circulating influenza virus strains tested by the WHO have been susceptible to neuraminidase inhibitors
- ▶ Reduced susceptibility of some influenza virus strains, particularly influenza A(H1N1), to oseltamivir or peramivir can emerge during or after treatment, especially in young children and immunocompromised patients with prolonged viral shedding
- ▶ Resistant isolates have usually remained susceptible to zanamivir, but reduced susceptibility to zanamivir has been reported
- ▶ Baloxavir is not recommended for severely immunosuppressed patients because of concerns about emergence of resistance

### References:

1. Approximate WAC for 5 days' treatment with oseltamivir capsules or zanamivir, or for a single treatment dose of peramivir or baloxavir, at the usual adult dosage. WAC = wholesaler acquisition cost, or manufacturer's published price to wholesalers; WAC represents published catalogue or list prices and may not represent an actual transactional price. Source: Analysource® Monthly. November 5, 2022. Reprinted with permission by First Databank, Inc. All rights reserved. ©2022. [www.fdbhealth.com/policies/drug-pricing-policy](http://www.fdbhealth.com/policies/drug-pricing-policy).
2. CDC. Influenza antiviral medications: summary for clinicians. Available at: <https://www.cdc.gov/flu/professionals/antivirals/summary-clinicians.htm>. Accessed November 5, 2022.
3. Antiviral drugs for influenza for 2022-2023. *Med Lett Drugs Ther* 2022; 64:185.
4. Although not FDA-approved for use in children <2 weeks old, the CDC recommends children <2 weeks old be treated with 3 mg/kg bid. For treatment of premature infants, refer to CDC recommendations ([www.cdc.gov/flu](http://www.cdc.gov/flu)).
5. The American Academy of Pediatrics has recommended a dose of 3.5 mg/kg for infants 9-11 months old based on the results of a study showing that a higher dose was needed to achieve the target exposure in this age group (DW Kimberlin et al. *J Infect Dis* 2013;207:709).
6. Although not FDA-approved for prophylaxis in children <1 year old, the American Academy of Pediatrics and CDC recommend that children 3 months-<1 year old receive 3 mg/kg once/day. Chemoprophylaxis is generally not recommended for premature infants or infants <3 months old.

**PRESIDENT:** Mark Abramowicz, M.D.; **VICE PRESIDENT, EDITOR IN CHIEF:** Jean-Marie Pflomm, Pharm.D.; **ASSOCIATE EDITORS:** Susan M. Daron, Pharm.D., Amy Faucard, MLS, Michael P. Viscusi, Pharm.D. **CONSULTING EDITORS:** Joanna Esterow, PA-C, Mordechai Sacks, DMSc, PA-C, Brinda M. Shah, Pharm.D., F. Peter Swanson, M.D.

**CONTRIBUTING EDITORS:** Carl W. Bazil, M.D., Ph.D., Columbia University College of Physicians and Surgeons; Ericka L. Crouse, Pharm.D., B.C.P.P., C.G.P., F.A.S.H.P., F.A.S.C.P., Virginia Commonwealth University; Vanessa K. Dalton, M.D., M.P.H., University of Michigan Medical School; Eric J. Epstein, M.D., Albert Einstein College of Medicine; David N. Juurlink, BPhm, M.D., Ph.D., Sunnybrook Health Sciences Centre; Richard B. Kim, M.D., University of Western Ontario; Sandip K. Mukherjee, M.D., F.A.C.C., Yale School of Medicine; Dan M. Roden, M.D., Vanderbilt University School of Medicine; Esperance A.K. Schaefer, M.D., M.P.H., Harvard Medical School; Neal H. Steigbigel, M.D., New York University School of Medicine; Arthur M. F. Yee, M.D., Ph.D., F.A.C.R., Weill Medical College of Cornell University

**MANAGING EDITOR AND DIRECTOR OF CONTENT OPERATIONS:** Susie Wong; **EDITORIAL ASSISTANT:** Karrie Ferrara

**FULFILLMENT AND SYSTEMS MANAGER:** Cristine Romatowski; **EXECUTIVE DIRECTOR OF SALES:** Elaine Reaney-Tomaselli

**EXECUTIVE DIRECTOR OF MARKETING AND COMMUNICATIONS:** Joanne F. Valentino; **INTERIM PUBLISHER:** Jean-Marie Pflomm, Pharm.D.

Founded in 1959 by Arthur Kallet and Harold Aaron, M.D.

**Copyright and Disclaimer:** The Medical Letter, Inc. is an independent nonprofit organization that provides healthcare professionals with unbiased drug prescribing recommendations. The editorial process used for its publications relies on a review of published and unpublished literature, with an emphasis on controlled clinical trials, and on the opinions of its consultants. The Medical Letter, Inc. does not sell advertising or receive any commercial support. No part of the material may be reproduced or transmitted by any process in whole or in part without prior permission in writing. The Medical Letter, Inc. does not warrant that all the material in this publication is accurate and complete in every respect. The Medical Letter, Inc. and its editors shall not be held responsible for any damage resulting from any error, inaccuracy, or omission.

#### Subscription Services

**Address:**  
The Medical Letter, Inc.  
145 Huguenot St. Ste. 312  
New Rochelle, NY 10801-7537  
www.medicalletter.org

**Customer Service:**  
Call: 800-211-2769 or 914-235-0500  
Fax: 914-632-1733  
E-mail: [custserv@medicalletter.org](mailto:custserv@medicalletter.org)

**Permissions:**  
To reproduce any portion of this issue,  
please e-mail your request to:  
[permissions@medicalletter.org](mailto:permissions@medicalletter.org)

**Subscriptions (US):**  
1 year - \$159; 2 years - \$298;  
3 years - \$398. \$65 per year  
for students, interns, residents,  
and fellows in the US and Canada.  
Reprints - \$45 per issue or article

**Site License Inquiries:**  
E-mail: [SubQuote@medicalletter.org](mailto:SubQuote@medicalletter.org)  
Call: 800-211-2769  
Special rates available for bulk  
subscriptions.

Get Connected:    

Copyright 2022. ISSN 0025-2859

The  
Medical  
Letter